



## ig® Stable 2 Chemically Competent Cells

### Manual

|                     |                |                |
|---------------------|----------------|----------------|
| <b>Catalog #</b>    | <b>1016-12</b> | <b>1016-24</b> |
| <b>Package Size</b> | 12x50µl        | 24x50µl        |



### Important!

#### **-80°C Storage Required**

- \* Immediately inspect packages
- \* Freeze upon receipt



visit us online for more  
products & custom services

**Intact Genomics, Inc.**

## Table of Contents

|  |          |
|--|----------|
| <b>Product Description.....</b>                      | <b>3</b> |
| <b>Specifications.....</b>                           | <b>3</b> |
| <b>Reagents Needed for One Reaction.....</b>         | <b>3</b> |
| <b>Product Components &amp; Storage.....</b>         | <b>3</b> |
| <b>Genotype.....</b>                                 | <b>3</b> |
| <b>Product Benefits.....</b>                         | <b>4</b> |
| <b>Quality Control.....</b>                          | <b>4</b> |
| <b>General Guidelines.....</b>                       | <b>4</b> |
| <b>Calculation of Transformation Efficiency.....</b> | <b>5</b> |
| <b>High Efficiency Transformation Protocol.....</b>  | <b>5</b> |
| <b>5 Minute Transformation Protocol.....</b>         | <b>6</b> |
| <b>Related Products.....</b>                         | <b>6</b> |
| <b>Ordering Information.....</b>                     | <b>6</b> |
| <b>Technical Support.....</b>                        | <b>7</b> |

### Description:

Intact Genomics (ig<sup>®</sup>) Stable 2 chemically competent *E. coli* cells are suitable for high efficiency transformation in a wide variety of applications such as cloning and sub-cloning. Stable 2 cells are capable of cloning methylated genomic sequences, retroviral sequences and direct repeat sequences. Intact Genomics Stable 2 cells provide superb transformation efficiency, allowing for increased opportunity for experimental success.

### Specifications:

**Competent cell type:** Chemically Competent

**Species:** *E. coli*

**Format:** Tubes

**Transformation efficiency:**  $\geq 1.0 \times 10^9$  cfu/ $\mu$ g pUC19 DNA

**Blue/white screening:** No

**Shipping condition:** Dry ice

### Reagents Needed for One Reaction:

- ig<sup>®</sup> Stable 2 chemically competent cells: 50  $\mu$ l
- DNA (or pUC19 Control, 10 pg/ $\mu$ l): 1  $\mu$ l
- Recovery medium: 1 ml

### Product Components & Storage:

- ig<sup>®</sup> Stable 2 competent cells: -80 °C
- pUC19 control DNA: -20 °C
- Recovery medium: 4 °C

### Genotype:

F- mcrA  $\Delta$ (mcrBC-hsdRMS-mrr) recA1 endA1lon gyrA96 thi supE44 relA1  $\lambda$ -  $\Delta$ (lac-proAB)

## Product Benefits:

ig<sup>®</sup> Stable 2 chemically competent cells have the following features:

- Stable 2 allows for cloning of methylated genomic sequences
- Stabilizes retroviral and direct repeat sequences including HIV
- High transformation efficiency allows aids in cloning rare sequences
- May be used for plasmids > 20 kb
- endA1 mutation increases plasmid yield significantly

## Quality Control:

Transformation efficiency is tested by using the pUC19 control DNA supplied with the kit and the high efficiency transformation protocol listed below. Transformation efficiency should be  $\geq 1 \times 10^9$  CFU/ $\mu$ g pUC19 DNA. Untransformed cells are tested for appropriate antibiotic sensitivity.

## General Guidelines:

Follow these guidelines when using ig<sup>®</sup> Stable 2 chemically competent *E. coli*.

- Handle competent cells gently as they are highly sensitive to changes in temperature or mechanical lysis caused by pipetting.
- Thaw competent cells on ice and transform cells immediately following thawing. After adding DNA, mix by tapping the tube gently. Do not mix cells by pipetting or vortexing.

## Example Calculation of TE:

Transformation Efficiency (TE) is defined as the number of colony forming units (cfu) produced by transforming 1µg of plasmid into a given volume of competent cells.

$$\text{TE} = \text{Colonies}/\mu\text{g}/\text{Dilution}$$

Transform 1 µl of (10 pg/µl) pUC19 control plasmid into 50 µl of cells, add 950 µl of Recovery Medium. Dilute 10 µl of this in 90 µl of Recovery Medium and plate 100 µl. Count the colonies on the plate the next day. If you count 200 colonies, the TE is calculated as follows:

$$\text{Colonies} = 200$$

$$\mu\text{g of DNA} = 0.00001$$

$$\text{Dilution} = 10/1000$$

$$\text{TE} = 200/.00001/0.01 = 2.0 \times 10^9$$

## High Efficiency Transformation Protocol:

Use this procedure to transform ig<sup>®</sup> Stable 2 chemically competent cells. We recommend verifying the transformation efficiency of the cells using the pUC19 control DNA supplied with the kit. Do not use these cells for electroporation.

- 1) Remove competent cells from the -80 °C freezer and thaw completely on wet ice (10-15 minutes).
- 2) Aliquot 1-5 µl (1 pg-100 ng) of DNA to the chilled microcentrifuge tubes on ice.
- 3) When the cells are thawed, add 50 µl of cells to each DNA tube on ice and mix gently by tapping 4-5 times. For the pUC19 control, add 1 µl of (10 pg/µl) DNA to a chilled microcentrifuge tube, prior to adding 50 µl of cells. Mix well by tapping. Do not pipette up and down or vortex to mix, this can harm cells and decrease transformation efficiency.
- 4) Incubate the cells with DNA on ice for 30 minutes.
- 5) After 30 minute ice incubation, heat shock the cells at 42 °C for 45 seconds.
- 6) Transfer the tubes to ice for 2 minutes.
- 7) Add 950 µl of Recovery Medium or any other medium of choice to each tube.
- 8) Incubate tubes at 37 °C for 1 hour at 210 rpm.
- 9) Spread 50 µl to 200 µl from each transformation on Pre-warmed selection plates. We recommend plating two different volumes to ensure that at least one plate will have well-spaced colonies. For the pUC19 control, plate 100 µl on an LB plate containing 100 µg/ml ampicillin. Use sterilized spreader or autoclaved ColiRoller™ plating beads to spread evenly.
- 10) Incubate the plates overnight at 37 °C.

## 5 Minute Transformation Protocol:

The following procedure results in only ~10% of the transformation efficiency as the protocol listed above.

1. Remove competent cells from the -80 °C freezer and thaw in your hand.
2. Aliquot 1-5 µl (1 pg-100 ng) of DNA to the microcentrifuge tubes. Do not pipette up and down or vortex to mix, this can harm cells and decrease transformation efficiency.
3. Incubate the cells with DNA on ice for 2 minutes.
4. After 2 minute ice incubation, heat shock the cells at 42 °C for 45 seconds.
5. Transfer the tubes to ice for 2 minutes.
6. Add 950 µl of Recovery Medium at room temperature or any other medium of choice to each tube. Immediately spread 50 µl to 200 µl from each transformation on pre-warmed selection plates. We recommend plating two different volumes to ensure that at least one plate will have well-spaced colonies. For the pUC19 control, plate 50 µl on an LB plate containing 100 µg/ml ampicillin. Use sterilized spreader or autoclaved ColiRoller™ plating beads to spread evenly.
7. Incubate the plates overnight at 37 °C

## Related Products:

- ig<sup>®</sup> 5-Alpha Chemically Comp. Cells (Cat.# 1031-12)
- T4 DNA Ligase (Cat.# 3212)
- i7<sup>®</sup> High Fidelity DNA Polymerase (Cat.# 3254)
- Quick10™ Cloning Kit (Cat.# 4122)

## Ordering Information:

- Order online within the USA. Place orders on **www.intactgenomics.com** using our secure Shopping Cart.
- Order by email, phone, or fax.  
Email: **sales@intactgenomics.com**  
Phone: (314) 942-3655 | Toll-free : 855-835-7172 | Fax: (314) 942-3656
- Order via our distributors.

Intact Genomics owns the following registered trademarks granted by the United States Patent and Trademark Office (USPTO): Intact Genomics®, IG®, ig®, igTherapeutics®, FastAmp®, i7®, DirectPlate®.

All technology protocols discussed within this manual are assumed proprietary to Intact Genomics. This Product may be covered by pending or issued patents or may have certain limitations. Please contact us for more information. Purchase of this material conveys to buyer the non-transferable right to use the material purchased in research conducted by buyer, whether for teaching, non-commercial or commercial research purposes. Buyer may not sell or otherwise transfer these materials, its components, or unmodified descendants to a third party.

## Product Use Limitation and Disclaimers

This product is for research purposes only. It is not intended for therapeutic or diagnostic purposes in humans or animals. This product contains chemicals which may be harmful if misused or direct human contact is made.

Intact Genomics is dedicated to practicing and maintaining science and technology ethics. Buyer agrees to use the purchased materials in full compliance with applicable law and regulations.

## Technical Support & Customer Services

Intact Genomics (IG®) is dedicated to customer satisfaction regarding the use of our products for your research needs. Each new lot of our products is thoroughly tested to ensure it meets high quality standards and provides excellent results. We appreciate your business and your feedback regarding the performance of our products in your applications. Please follow the instructions carefully and contact us if additional assistance is needed.

Our hours are Monday - Friday, 8AM to 5PM, U.S. Central Standard Time.

### Intact Genomics, Inc.

11840 Westline Industrial Drive, Suite 120,  
St. Louis, MO. 63146, USA

**Phone:** (314) 942-3655 | **Toll-free :** 855-835-7172 | **Fax:** (314) 942-3656

**Email:** sales@intactgenomics.com | ig@intactgenomics.com

**Website:** www.intactgenomics.com

© 2024 Intact Genomics, Inc  
All Rights Reserved

