



igScript™ Glycerol-free Reverse Transcriptase

Manual

Catalog #	3342GF	3344GF
Package Size	10,000 units	50,000 units
Volume	50µl	250 µl
Concentration	200 units/µl	200 units/µl



Important!

-80°C Storage Required

- * Immediately inspect packages
- * Freeze upon receipt



visit us online for more
products & custom services

Intact Genomics, Inc.

Table of Contents

Product Description.....	3
Protein Purity.....	3
Product Source.....	3
Applications.....	3
Benefits.....	3
Product Components.....	3
Reaction Buffer.....	4
Storage.....	4
Unit Definition.....	4
Heat Inactivation.....	4
Synthesis Protocol.....	5
Related Products.....	6
Ordering Information.....	6
Technical Support	7

Description:

igScript™ Glycerol-free Reverse Transcriptase (igScript™ RT) is a recombinant Moloney Murine Leukemia Virus (MMLV) reverse transcriptase with reduced RNase H activity. It is highly efficient at producing full-length cDNA from long RNA templates. igScript™ RT is an RNA-directed DNA polymerase which lacks 3' → 5' exonuclease activity and is capable of producing cDNA from as little as 50 pg of total RNA for real-time RT-PCR analysis and other applications.

Protein Purity:

The physical purity of this enzyme is ≥98% as assessed by SDS-PAGE with Coomassie® blue staining.

Product Source:

The gene encoding recombinant MMLV Reverse Transcriptase with expressed in *E. coli*.

Applications:

- First strand cDNA synthesis for PCR or RT-PCR
- Gene expression data validation by using RT-PCR product

Benefits:

- Recombinant MMLV reverse transcriptase with greatly reduced RNase H activity.
- Active at temperatures up to 55°C.
- Highly efficient at producing full-length cDNA from as little as 50 pg of total RNA.
- Glycerol-free formulation for sensitive workflows

Product Components:

- igScript™ Reverse Transcriptase, Glycerol-free
- 10x igScript™ RT Reaction Buffer

1x MMLV Reverse Transcriptase Reaction Buffer:

- 50 mM Tris-HCl
- 75 mM KCl
- 3 mM MgCl₂
- 10 mM DTT
- pH 8.3 @ 25°C

Storage Temperature:

-80°C

Storage Buffer:

50 mM HEPES, 750 mM KCl, 1 mM DTT, 0.1 mM EDTA, pH 7.5 @ 25°C

Unit Definition:

One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into acid insoluble material in 10 minutes at 37 °C using poly r(A)/oligo (dT) as a substrate.

Heat Inactivation:

65°C for 20 minutes.

First Strand cDNA Synthesis Protocol:

- 1) In a sterile micro-centrifuge tube, add the following components on ice:

Component	Volume
Total RNA	Up to 1.0 µg
Primer: d(T) ₂₃ VN (50 µM) or Random Primer Mix (60 µM) or Gene Specific Primer (10 µM)	2.0 µl
10 mM dNTPs	1.0 µl
H ₂ O	Up to 10.0 µl

- 2) Heat the reaction for 5 minutes at 65°C. Spin down briefly. Place the tube immediately on ice.

- 3) Add the following components:

Component	Volume
10x igScript™ RT Reaction Buffer	2.0 µl
igScript™ Reverse Transcriptase (200 U/µl)	1.0 µl
RNase Inhibitor (40 U/µl)*	0.2 µl
H ₂ O	6.8 µl

- 4) If using random hexamers, incubate the reaction at 25° C for 10 minutes, then proceed to step 5.
- 5) Incubate the reaction at 42°C for 30-60 minutes.
- 6) Inactivate the enzyme by incubating at 65°C for 20 minutes.
- 7) Store products at -20°C or proceed to next step.

* RNase Inhibitor, Murine (with glycerol or without, Cat.# 3714 or 3714GF)

Related Products:

- RNase Inhibitor, Murine (Cat.# 3714)
- Glycerol-free RNase Inhibitor, Murine (Cat.# 3714GF)
- igScript™ One Step RT-qPCR Kit (Cat.# 4214)
- igScript™ First Strand cDNA Synthesis Kit (Cat.# 4312)
- ig® SYBR Green qPCR 2x Master Mix (Cat.# 3354)
- igScript™ Probe Based One Step RT-qPCR Kit (Cat.#4243, 4245, 4247)

Ordering Information:

- Order online within the USA. Place orders on **www.intactgenomics.com** using our secure Shopping Cart.
- Order by email, phone, or fax.
Email: **sales@intactgenomics.com**
Phone: (314) 942-3655 | Toll-free : 855-835-7172 | Fax: (314) 942-3656
- Order via our distributors.

Intact Genomics owns the following registered trademarks granted by the United States Patent and Trademark Office (USPTO): Intact Genomics®, IG®, ig®, igTherapeutics®, FastAmp®, i7®, DirectPlate®.

All technology protocols discussed within this manual are assumed proprietary to Intact Genomics. This Product may be covered by pending or issued patents or may have certain limitations. Please contact us for more information. Purchase of this material conveys to buyer the non-transferable right to use the material purchased in research conducted by buyer, whether for teaching, non-commercial or commercial research purposes. Buyer may not sell or otherwise transfer these materials, its components, or unmodified descendants to a third party.

Product Use Limitation and Disclaimers

This product is for research purposes only. It is not intended for therapeutic or diagnostic purposes in humans or animals. This product contains chemicals which may be harmful if misused or direct human contact is made.

Intact Genomics is dedicated to practicing and maintaining science and technology ethics. Buyer agrees to use the purchased materials in full compliance with applicable law and regulations.

Technical Support & Customer Services

Intact Genomics (IG®) is dedicated to customer satisfaction regarding the use of our products for your research needs. Each new lot of our products is thoroughly tested to ensure it meets high quality standards and provides excellent results. We appreciate your business and your feedback regarding the performance of our products in your applications. Please follow the instructions carefully and contact us if additional assistance is needed.

Our hours are Monday - Friday, 8AM to 5PM, U.S. Central Standard Time.

Intact Genomics, Inc.

11840 Westline Industrial Drive, Suite 120,
St. Louis, MO. 63146, USA

Phone: (314) 942-3655 | **Toll-free :** 855-835-7172 | **Fax:** (314) 942-3656

Email: sales@intactgenomics.com | ig@intactgenomics.com

Website: www.intactgenomics.com

© 2024 Intact Genomics, Inc
All Rights Reserved

