


T4 UvsY Protein

Manual



Important!

-20°C Storage Required

- * Immediately inspect packages
- * Freeze upon receipt

FOR RESEARCH USE ONLY. NOT FOR HUMAN OR DIAGNOSTIC USE

Catalog #	3572	3575	3577
Package Size	100 µg	500 µg	1000µg

Intact Genomics, Inc.

www.intactgenomics.com



Description:

Intact Genomics (ig®) UvsY is the phage T4 recombination mediator protein, and structural and biophysical studies provide insights into its role in T4 homologous recombination. During T4 homologous recombination, the UvsX recombinase must compete with the prebound gp32 single-stranded binding protein for DNA-binding sites and UvsY stimulates this filament nucleation event (1). UvsY is a 15.8-kDa protein with properties that are consistent with its role as mediator; it stimulates the DNA-dependent ATPase activity of UvsX, lowers the critical concentration of UvsX that is required for activity, and promotes strand exchange (2, 3). UvsY efficiently promotes the UvsX-catalyzed strand invasion reaction by recruiting to ssDNA–gp32 complexes, promotes the release of gp32, and favors the binding of UvsX.

Protein Purity:

The physical purity of this enzyme is ≥98% as assessed by SDS-PAGE with Coomassie® blue staining (Fig. 1).

Product Source:

E. coli BL21 (DE3) strain expressing T4 UvsY gene with an N-terminal His tag.

This GST tag does not have any effect on UvsY activity.

Product Includes:

- UvsY protein
- 10x UvsY Reaction Buffer

1x UvsY Reaction Buffer:

- 20 mM Tris-acetate, 100 mM Potassium acetate
- 10 mM Magnesium acetate, 1 mM DT
- pH 7.8 @ 25°C

Storage Buffer:

50 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA

50% Glycerol, pH 7.5 @ 25°C




Fig. 1:
Lane 1, Protein marker
Lane 2, UvsY.

Quality Control:

UvsY is free from detectable nuclease activities.

References:

1. Beernink HT, Morrical SW. RMPs: Recombination/replication mediator proteins. Trends Biochem Sci. 1999;24(10):385–389.
2. Hashimoto K, Yonesaki T. The characterization of a complex of three bacteriophage T4 recombination proteins, uvsX protein, uvsY protein, and gene 32 protein, on single-stranded DNA. J Biol Chem. 1991;266(8):4883–4888.
3. Beernink HT, Morrical SW. The uvsY recombination protein of bacteriophage T4 forms hexamers in the presence and absence of single-stranded DNA. Biochemistry. 1998;37(16):5673–5681

Related Products:

1. T4 gp32 Protein (Cat.# 3515)
2. T4 UvsX Protein (Cat.# 3562,)
3. Bsu DNA Polymerase (Cat.# 3585)
4. Sau DNA Polymerase (Cat.# 3595)
5. Exonuclease III (Cat.# 3415)
6. Exonuclease IV (Nfo) (Cat.# 3425)

Technical Support:

Intact Genomics is committed to supporting the worldwide scientific research community by supplying the highest quality reagents. Each new lot of our products is tested to ensure they meet the quality standards and specifications designated for the product.

Please follow the instructions carefully and contact us if additional assistance is needed. We appreciate your business and your feedback regarding the performance of our products in your applications.

Enzyme Concentration:

IG uses orthogonal, 3-part approaches to determine the enzyme concentration to provide you with consistent and reliable enzymes for your needs. The quantity of a protein sample is assessed using densitometry with polyacrylamide gel electrophoresis (PAGE), UV absorbance spectra of native protein, and using a protein standard assay such as bicinchoninic acid assay (BCA) using bovine serum albumin (BSA) as a standard (Figure 1).

Why does IG use all three approaches?

1. Each method above has limitations. The limitations include experimental noise, accuracy, and susceptibility to buffer and/or enzyme conditions.
2. Each enzyme has unique physical properties that make a single approach to analyzing proteins a challenge. Each enzyme has a different protein sequence, different requirements to be stable in solution, and different requirements to retain its maximal activity. These differences can interfere with or convolute results, especially when compared to other enzymes. When used together, however, each method provides the scientist with independent measures of both enzyme and buffer purity and quality.




Figure 1: Enzyme quantitation methods used by IG. A) SDS-polyacrylamide gel electrophoresis. Ladder in 1st lane, 2 μ g BSA (~67 kDa) as a standard in lanes 1-3, and IG enzymes (~40 kDa) in lanes 4-6. The yellow boxes are the areas evaluated by densitometry. The integrated band intensities of IG enzymes are compared with integrated band intensities from BSA to assay concentration. B) UV spectrum of a clean IG enzyme with protein peaks at 230 nm and at 280 nm. An extinction coefficient at 280 nm is typically used to quantify protein using these spectra with buffer subtraction at 330 nm. C) BCA standard curve for BSA. The curve is used to calculate an IG enzyme concentration using BSA as the standard.

Intact Genomics is committed to supporting the worldwide scientific research community by supplying the highest quality life science products to help research scientists worldwide discover solutions to critical challenges in human health, agriculture and the environment. Each new lot of our products is tested to ensure they meet the quality standards and specifications designated for the product. Please follow the instructions carefully and contact us if additional assistance is needed.

We appreciate your business and your feedback regarding the performance of our products in your applications. Our hours are Monday - Friday, 8AM to 5PM, U.S. central standard time.

<http://www.intactgenomics.com>

Intact Genomics, Inc.
11840 Westline Industrial Drive, Suite 120
St. Louis, MO. 63146

Phone: (314) 942-3655 | **Toll-free :** 855-835-7172 | **Fax:** (314) 942-3656

Email: sales@intactgenomics.com

Legal Notices:

Intact Genomics owns the following registered trademarks granted by the United States Patent and Trademark Office (USPTO): Intact Genomics®, IG®, ig®, igTherapeutics®, FastAmp®, i7®, DirectPlate®.

All technology protocols discussed within this manual are assumed proprietary to IG's products, and if registered, are protected by US patent and trademark laws.

Notice to Buyer - Limited License:

Buyer agrees by purchasing the product that it's intended use is for research purposes only, not for human or diagnostic use. Purchase of this material conveys to buyer the non-transferable right to use the material purchased in research conducted by buyer, whether for teaching, non-commercial or commercial research purposes. Buyer may not sell or otherwise transfer these materials, its components, or unmodified descendants to a third party. Use of these products maybe covered by IG's patents or trademarks.

Products are understood by buyer to be experimental in nature and may have hazardous properties. Unless prohibited by law, buyer assumes all liability for claims or damages against it by third parties that relate to or arise from the use, storage, or disposal of the purchased materials.

Buyer agrees to use the purchased materials in full compliance with applicable law and regulations. Intact Genomics will not be held liable for activities outside of itself and the products intended use.

For more information or to express legal concerns relating to patent and trademarks, please contact us at: **info@intactgenomics.com** or **sales@intactgenomics.com**

Thank you for your business!

www.intactgenomics.com