



igSuper™ Tte UvrD Helicase

Manual

Catalog #	3717	3719
Package Size	2 µg	40 µg
Concentration	20 µg/mL	20 µg/mL
Volume	100 µL	2 mL



Important!

-20°C Storage Required

- * Immediately inspect packages
- * Freeze upon receipt



visit us online for more
products & custom services

Intact Genomics, Inc.

Table of Contents

Product Description..... 3

Applications..... 3

Benefits..... 3

Source..... 3

Specifications..... 3

Components and Storage Temperature..... 4

Quality Control..... 4

Protocol 4

References..... 5

Related Products 5

Ordering Information..... 5

Technical Support and Customer Service 6

Description:

Intact Genomics igSuper™ Tte UvrD is a thermostable, ATP-dependent DNA helicase derived from the thermophilic bacterium *Thermoanaerobacter tengcongensis* (Tte). The enzyme unwinds double-stranded DNA (dsDNA), generating single-stranded DNA (ssDNA) that can be accessed by other proteins, enzymes, primers, and probes.

igSuper™ Tte UvrD is an engineered version of Tte UvrD for potential improved characters in literatures¹ and a key enzyme used in helicase-dependent amplification (HDA)². In HDA, igSuper™ Tte UvrD separates complementary DNA strands to generate single-stranded templates, allowing primer hybridization and DNA polymerase extension without the thermal cycling required for PCR. Studies have also shown that HDA may use the same primer sequences designed for PCR, helping simplify assay development³.

igSuper™ Tte UvrD is thermostable and it is active at temperatures up to 65 °C. It is suitable for HDA, DNA-unwinding studies, helicase activity assays, isothermal nucleic acid amplification, and other molecular biology applications requiring enzymatic strand separation.

Applications:

- Helicase-dependent amplification (HDA)
- Isothermal nucleic acid amplification
- DNA strand-unwinding studies
- Helicase activity assays
- Probe- and primer-based DNA detection workflows

Benefits:

- Thermostable and active at temperatures up to 65 °C
- Generates single-stranded DNA without thermal cycling
- Supports primer hybridization and DNA polymerase extension
- May be used with PCR-designed primers, simplifying assay development
- Suitable for research and diagnostic assay development

Source:

Recombinant UvrD expressed from a plasmid carrying the engineered thermostable UvrD gene from *Thermoanaerobacter tengcongensis*.

Specifications:

- Purified recombinant igSuper™ Tte UvrD Helicase
- Molecular weight: 82 kDa
- Free of detectable endonuclease activity

Product Components and Storage Temperature:

IG Cat #	Package Size	Component Name	IG Part #	Amount	Concentration	Storage
3517	2 µg	igSuper™ Tte UvrD Helicase	P241281	1 × 100 µL	20 µg/mL	-20 °C
		10X Helicase Reaction Buffer	P241266	1 × 1 mL	10X	-20 °C
		10 mM Adenosine Triphosphate (ATP)	P241270	1 × 50 µL	10 mM	-20 °C
3519	40 µg	igSuper™ Tte UvrD Helicase	P241284	2 × 1 mL	20 µg/mL	-20 °C
		10X Helicase Reaction Buffer	P241266	1 × 1 mL	10X	-20 °C
		10 mM Adenosine Triphosphate (ATP)	P241274	1 × 200 µL	10 mM	-20 °C

Quality Control:

Nuclease Activity: DNA integrity is evaluated after overnight incubation with concentrated enzyme. The DNA must remain intact, with no detectable degradation, compared with a no-enzyme control.

Enzyme Activity: At 24 nM (2 ng/µL), the enzyme must pass both ATPase and fluorescence-based helicase activity assays and demonstrate performance comparable to an approved UvrD batch-control standard.

Protocol for DNA Helicase Activity:

The igSuper™ UvrD reaction measures ATP-dependent unwinding of complementary DNA strands.

- Set up the reaction according to the table below.

Reagent	µL of Sample
Template dsDNA	X (up to 1 µg)
Nuclease-Free H ₂ O	15-X
10x Helicase Buffer	2
10 mM ATP	2
igSuper Tte UvrD Helicase	1

Total Volume: 20 µL

- Incubate the reaction mixture at 65 °C for 20 minutes. The incubation time and temperature may be adjusted for different applications. Typical conditions range from 42–68 °C for 10–60 minutes. Note that ATP hydrolysis can limit helicase activity at higher temperatures.
- To inactivate the enzyme, incubate the reaction at 80 °C for 5 minutes.

References:

- Gavrilov, M., *et al.* Engineered helicase replaces thermocycler in DNA amplification while retaining desired PCR characteristics. *Nat Commun* 13, 6312 (2022).
- An, L. *et al.* Characterization of a Thermostable UvrD Helicase and Its Participation in Helicase-dependent Amplification *. *Journal of Biological Chemistry* 280, 28952–28958 (2005).
- Gavrilov, M. *et al.* Engineered helicase replaces thermocycler in DNA amplification while retaining desired PCR characteristics. *Nat Commun* 13, 6312 (2022).

Related Products:

- IG® Tte UvrD Helicase (Cat.# 3711, 3713)
- IG® Tte MutL Protein (Cat.# 3722, 3724)
- IG® RPA Master Mix (Cat.# 3545, 3547)
- IG® RPA Kit v2 (Cat.# 3533, 3534)
- T4 UvsX DNA Recombinase (Cat.# 3562, 3565)
- T4 UvsY Protein (Cat.# 3572, 3575)

Ordering Information:

- Order online within the USA. Place orders on **www.intactgenomics.com** using our secure Shopping Cart.
- Order by email, phone, or fax.
Email: **sales@intactgenomics.com**
Phone: (314) 942-3655 | Toll-free : 855-835-7172 | Fax: (314) 942-3656
- Order via our distributors.



Intact Genomics owns the following registered trademarks granted by the United States Patent and Trademark Office (USPTO): Intact Genomics®, IG®, ig®, FastAmp®, i7®, DirectPlate®.

All technology protocols discussed within this manual are assumed proprietary to Intact Genomics. This Product may be covered by pending or issued patents or may have certain limitations. Please contact us for more information.

Product Use Limitation and Disclaimers

This product is intended primarily for research purposes. By purchasing this material, the buyer is granted a non-transferable license to use it in research conducted by the buyer, including teaching, non-commercial, or commercial research activities. The buyer may not sell, distribute, or otherwise transfer this material, any of its components, or any unmodified derivatives to any third party without prior written consent from Intact Genomics.

Intact Genomics is dedicated to practicing and maintaining science and technology ethics. Buyer agrees to use the purchased materials in full compliance with applicable law and regulations.

Technical Support & Customer Services

Intact Genomics (IG®) is dedicated to customer satisfaction regarding the use of our products for your research needs. Each new lot of our products is thoroughly tested to ensure it meets the high quality standards and provide excellent results. We appreciate your business and your feedback regarding the performance of our products in your applications. Please follow the instructions carefully and contact us if additional assistance is needed.

Our hours are Monday - Friday, 8 AM to 5 PM, U.S. central standard time (CST).

Intact Genomics, Inc.

11840 Westline Industrial Drive, Suite 120,
St. Louis, MO. 63146, USA

Phone: (314) 942-3655 | **Toll-free :** 855-835-7172 | **Fax:** (314) 942-3656

Email: sales@intactgenomics.com | ig@intactgenomics.com

Website: www.intactgenomics.com

© 2026 Intact Genomics, Inc
All Rights Reserved

